

SYMBIODINIUM (PYRRHOPHYTA) GENOME SIZES (DNA CONTENT) ARE SMALLEST AMONG DINOFLAGELLATES¹

*Todd C. LaJeunesse*²

Department of Biology, Florida International University, Miami, Florida 33199 USA

Georgina Lambert

Department of Plant Sciences, University of Arizona, Tucson, Arizona 85721-0036 USA

Robert A. Andersen

Provasoli-Guillard National Center for Culture of Marine Phytoplankton, Bigelow Laboratory for Ocean Sciences, West Boothbay Harbor, Maine 04575 USA

Mary Alice Coffroth

Department of Biological Sciences, State University of New York, Buffalo, New York 14260 USA

and

David W. Galbraith

Department of Plant Sciences, University of Arizona, Tucson, Arizona 85721-0036 USA

Using flow cytometric analysis of fluorescence, we measured the genome sizes of 18 cultured “free-living” species and 29 *Symbiodinium* spp. isolates cultured from stony corals, gorgonians, anemones, jellyfish, and giant clams. Genome size directly correlated with cell size, as documented previously for most eukaryotic cell lines. Among the smallest of dinoflagellates, *Symbiodinium* spp. (6–15 μm) possessed the lowest DNA content that we measured (1.5–4.8 $\text{pg} \cdot \text{cell}^{-1}$). Bloom-forming or potentially harmful species in the genera *Alexandrium*, *Karenia*, *Pfiesteria*, and *Prorocentrum* possessed genomes approximately 2 to 50 times larger in size. A phylogenetic analysis indicated that genome/cell size has apparently increased and decreased repeatedly during the evolution of dinoflagellates. In contrast, genome sizes were relatively consistent across distantly and closely related *Symbiodinium* spp. This may be the product of intracellular host habitats imposing strong selective pressures that have restricted symbiont size.

Key index words: C-values; dinoflagellates; genome size; *Symbiodinium*; zooxanthellae

Abbreviation: DAPI, 4',6-diamidino-2-phenylindole; ITS, internal transcribed spacer

Dinoflagellates (Pyrrophyta) are primarily unicellular biflagellates. Collectively, they form one of three

main clades comprising the alveolates, an ancient eukaryotic lineage that includes the ciliates and apicomplexan parasites. Dinoflagellates occur in both marine and freshwater habitats and are common constituents of phytoplankton communities. Of the more than 2000 known species, some are responsible for harmful algal blooms, whereas other taxa are appreciated for their impressive bioluminescence. Each species exhibits either phototrophy (photosynthesis), heterotrophy, or mixotrophy (Taylor 1987). Their ecological breadth also extends to parasitic and mutually symbiotic relations involving numerous vertebrate and invertebrate hosts (Cachon 1987, Trench 1993). The widespread symbioses between *Symbiodinium* and coral reef-dwelling invertebrates are especially notable because these mutualistic associations are essential for the existence of coral reef ecosystems (Muscatine and Porter 1977, Muscatine 1990).

Scientific fascination with dinoflagellates extends far beyond their obvious ecological and economic importance. They possess numerous unique cellular, molecular, and biochemical traits not observed in “text book” life forms and model organisms. Such disparities include 1) permanently condensed chromosomes lacking conventional histones; 2) DNA that contains some hydroxymethyluracil in place of thymine, unique among eukaryotes (Rizzo 1987); 3) unique toxic molecules and sterols (Shimizu 1987, Withers 1987); 4) complex organelle structures such as light sensing ocelloids, ejectile nematocyst and trichocysts, and the pusule, an invagination probably involved in osmoregulation and nutrient acquisition (Dodge and Greuet 1987); 5) an elaborate cell covering or thecal system (amphiesma) (Dodge and Greuet 1987); and 6) un-

¹Received 16 December 2004. Accepted 30 April 2005.

²Author for correspondence: e-mail lajeunes@fiu.edu.

sual genes and gene arrangements, such as bacterial type II RUBISCO (Whitney et al. 1995) and minicircular plastid DNA (Zhang et al. 1999). Collectively, these and other unique cytological, ultrastructural, and genetic attributes present intriguing topics of research. In summary, examination of the dinoflagellates underscores the degree of divergence that has occurred in the evolution of eukaryotic life forms.

The economic, ecological, and evolutionary significance of dinoflagellates establishes a persuasive argument for sequencing genomes representative of this group. However, one major obstacle to achieving this goal is that dinoflagellates typically possess genomes many times greater than the $3.0 \text{ pg} \cdot \text{cell}^{-1}$ (approximately) of the human genome. For example, *Prorocentrum micans* has one of the largest genomes measured for photosynthetic protists (e.g. 230 to $280 \text{ pg} \cdot \text{cell}^{-1}$; Velduis et al. 1997). Because past surveys of genome size among dinoflagellates were taxonomically limited (Rizzo 1987, Velduis et al. 1997, Parrow and Burkholder 2002), species may yet exist with small genomes more suitable for sequencing.

For this primary reason, we surveyed genome sizes (C-values, defined as the DNA content per haploid nucleus) for a range of cultured dinoflagellates via flow-cytometric analysis of 4',6-diamidino-2-phenylindole (DAPI, Sigma Inc., St. Louis, MO, USA) stained cells for a range of cultured dinoflagellates. Our focus was on *Symbiodinium*, but we included representatives from familiar bloom-forming or potentially toxic genera (e.g. *Alexandrium*, *Karenia*, and *Pfiesteria*) as well as species from other genera available in culture. These were compared with measurements on various *Symbiodinium* spp. in the subgeneric clades A, B, C, D, E, and F (reviewed in Baker 2003). Sequences of the highly conserved (Maroteaux et al. 1985) and functionally constrained (Elela and Nazar 1997) 5.8S ribosomal gene (rDNA) provided the basis of a simple phylogenetic reconstruction involving most of the taxa examined in this study. Based on this phylogeny, we trace the evolution of genome/cell size across these lineages. Compared with other lineages, the evolutionarily divergent *Symbiodinium* spp. possess limited genome and cell sizes.

MATERIALS AND METHODS

Culturing and fixation. Cells were harvested while in exponential (log) growth to maintain consistent and accurate comparisons of nuclear DNA content (Allen et al. 1975, Loeblich 1976). After centrifugation ($1000g$ for 4 min), the resulting pellet (100,000–250,000 cells) was fixed with 1 to 10 mL of methanol:acetic acid (3:1 vol/vol) and then placed on ice for 20 min. The cells were recovered by centrifugation and the supernatant removed. The fixation step was repeated two or three times, with the addition of fresh fixative. This process removed essentially all the chloroplast pigments that might potentially interfere with subsequent nuclear fluorescence detection. After a fourth centrifugation and decantation, a 90% methanol wash was used to remove the acetic acid. After 5 min on ice, the cells were centrifuged and a fresh volume of 90% methanol was added. The fixed cells were

then immediately shipped at 4°C by overnight courier to the University of Arizona for DNA measurements.

DNA content measurements. Cells were recovered by centrifugation at $1000g$ for 4 min. They were stained by addition of 4 mL of a solution containing $20 \text{ g} \cdot \text{mL}^{-1}$ DAPI buffered by 20 M MOPS, pH 7. DNA content was measured using the DNA stain DAPI on a CCA-II Flow Cytometer (Partec, Munster, Germany) that was aligned using AlignFlo fluorescent microspheres (Molecular Probes, Eugene, OR, USA). Flow rates were typically $500 \text{ cells} \cdot \text{s}^{-1}$, and uniparametric histograms were accumulated to at least 1000 events. Coefficient of variation values for the DNA content peaks were typically less than 10%. DNA content measurements were calibrated using chicken red blood cells and *Arabidopsis thaliana* and *Nicotiana tabacum* nuclei (Johnston et al. 1999). Calculations were performed based on the chicken red blood cell standard of 3.0 pg per nucleus (Johnston et al. 1999), using a conversion factor of $1 \text{ pg} = 9.65 \times 10^8 \text{ bp}$ (Bennett 2000). To define the degree of technical variability in the method of measurement, the same culture of "*Gymnodinium*" *simplex* (CCMP419) was measured seven times on three different occasions (quotation marks designate taxonomic uncertainty [Daughbjerg et al. 2000] or that the taxon lacks a formal description). In addition, replicate samples for other taxa were analyzed whenever possible (see Results).

The asexual vegetative stages of all dinoflagellates (with the possible exception of *Noctiluca*) are thought to be haploid (haplontic; Pfiester and Anderson 1987, Coats 2002). For example, nuclear reconstructions (Blank 1987) and molecular genetic evidence (Santos and Coffroth 2003) indicate that *Symbiodinium* spp. are haploid. For the purposes of this study, we assumed that the cells of each species were haploid.

Fluorescence emission from DNA stained with DAPI is proportional to the adenine and thymine (AT) content. Major across-species differences in the proportion of these nucleotides relative to cytosine and guanine (GC) would affect our DNA measurements. The AT proportions previously measured for *Symbiodinium* sp. representatives in clades A, C, and F ranged between 50.0% and 53.6% (Blank and Huss 1989). The conservation in AT/GC ratios should not adversely impact conclusions drawn from measurements of *Symbiodinium* spp. C-values. However, for species such as *Heterocapsa pygmaea* whose nuclear AT content has been measured as 36%, we expect a systematic underestimation of C-values relative to previous findings (Triplett et al. 1993). The degree of underestimation should not affect conclusions regarding species suitable for genome sequencing because it would at worst represent an impossible factor of 2.

To test the relative precision of our measurements, C-value estimates made for several *Symbiodinium* spp. were compared against previously published estimates of chromosome volumes. In the dinokaryotic nucleus, chromosomes are condensed throughout mitosis and interphase. Therefore, chromosome volume was calculated from three-dimensional reconstructions based on transmission electron micrographs taken of serial sections through the nucleus (Blank and Trench 1985, Trench and Blank 1987, Blank and Huss 1989). Ideally, the use of isolated nuclei would have improved our accuracy because staining whole cells results in some nonspecific background fluorescence.

Cell size measurements. Light microscopy and an eye-piece reticule calibrated with a stage micrometer were used to measure cell sizes. Cell length ranges for the free-living species were obtained from the Center for Culture of Marine Phytoplankton (CCMP) database (Table 1). For *Symbiodinium* spp., coccoid cell sizes can vary considerably within a culture (Domotor and D'Elia 1986). To minimize the variability of this measurement, observations were made along the division plane of dividing cells (doublets) from cultures in log

TABLE 1. The DNA content in cultured dinoflagellates from non-*Symbiodinium* genera.

Taxon (Dinophyceae)	CCMP culture no.	Size in μm (median length)	$\text{pg} \cdot \text{cell}^{-1}$ (<i>n</i>)	Published flow cytometric and (nonflow cytometric) valuations in $\text{pg} \cdot \text{cell}^{-1}$	GenBank accession no.
<i>Alexandrium andersonii</i>	2222	20–24 (22)	21.8 (2)		
<i>Alexandrium insuetum</i>	2082	22–28 (25)	30.8 (2)		AB006996
<i>Alexandrium lusitanicum</i>	1888	17–19 (18)	31.0		AJ005050
<i>Alexandrium tamarense</i>	1598	20–30 (25)	103.5		AB006992
<i>Amphidinium carterae</i>	1314	12–18 (15)	5.9 (2)	9.8 ^a (3.2 ^b /2.7 ^c)	
<i>Gymnodinium simplex</i>	419	8–12 (10)	11.6 (7)		AY686651
<i>Heterocapsa triquetra</i>	449	18–20 (19)	24.1	19.1 ^a	AF527816
<i>Heterocapsa pygmaea</i>	1322	9–14 (11.5)	3.8 (2)	(9.10 ^d)	AB084094
<i>Polarella glacialis</i>	2088	9–14 (11.5)	7.0 (2)		
<i>Karenia brevis</i>	2229	20–28 (24)	57.1	(101.6 ^c /112.5 ^c)	AF352369
<i>Karenia mikimotoi</i>	429	28–30 (29)	100.1		AF318224
<i>Karlodinium galatheanum</i>	1974	10–18 (14)	16.9 (2)		AF172716
<i>Karlodinium galatheanum</i>	2282	8–20 (14)	16.3		AF352366
<i>Katodinium rotundatum</i>	1542	11–16 (13.5)	3.6		
<i>Pfiesteria piscicida</i>	1830	10–12 (11)	5.5 (2)	5.7 ^c	AY245693
<i>Pfiesteria shumwayae</i>	2357	10–18 (14)	19.8	14.7 ^c	AY245694
<i>Prorocentrum ballicum</i>	1260	10–15 (12.5)	8.3 (2)		
<i>Prorocentrum dentatum</i>	1517	14–17 (15.5)	6.6 (2)		
<i>Prorocentrum micans</i>	689	22–38 (30)	115.2; 225.0	217.1 ^a (42.0 ^c)	AF370879
<i>Prorocentrum minimum</i>	1329	7–16 (12.5)	6.9 (3; SD = 0.35)		AF208244

Ranges in length dimensions (and median length) are provided. Numbers in parentheses refer to replicate flow cytometer readings. Accession numbers used to reconstruct the 5.8S phylogeny of Fig. 3 are also listed.

^aFlow cytometric values (Veldhuis et al. 1997) using SYTOX stain.

^bDiaminobenzoic acid C-value determinations by Holm-Hansen (1969).

^cDiaminobenzoic acid C-value (Thomas and Farquhar 1978) determinations by Rizzo (1987).

^dDiaminobenzoic acid C-value determinations by Triplet et al. (1993).

^eFlow cytometric values (Parrow and Burkholder 2002) using SYTOX stain.

phase (Schoenberg and Trench 1980, Fitt 1985). Size was calculated from an average of 10 to 30 of these doublets. Because isolates possessing the same internal transcribed spacer (ITS) type are similar in size (LaJeunesse 2001), not all isolates were measured. Size measurements on the slightly smaller and rapidly moving motile phase of *Symbiodinium* spp. were not taken (see discussion below).

Phylogenetic analysis. Sequences of the 5.8S ribosomal gene for the cultures used in this study were acquired from GenBank, with the exception of "*Gymnodinium simplex*" (Tables 1 and 2). The complete ITS1, 5.8S, and ITS 2 sequence for "*G. simplex*" was obtained following the methods described by LaJeunesse (2001). Daughjerg et al. (2000) showed that CCMP419 was not closely related to the type species of *Gymnodinium*, and therefore it should be reclassified. "*Symbiodinium trenchii*" (unpublished results) corresponds to "type" D1a and occurs in hosts from the Indian, Pacific, and Atlantic oceans. A sequence alignment consisting of 168 characters (including indels) of 18 distinctive sequences was prepared for phylogenetic analysis. Because members of a genus sometimes have the same 5.8S sequence, redundancies were omitted, leaving one sequence in the alignment to stand for more than one species. The proto-alveolate parasite, *Prekinsus marinus* (AF150990), was used as the outgroup (Saldarriaga et al. 2003). Maximum parsimony and maximum likelihood methods for phylogenetic reconstruction were performed using the software package PAUP v 4.10b (Swofford 2000).

RESULTS AND DISCUSSION

Genome size relates to cell size. This study extends the species of dinoflagellates for which genome size measurements are available, most notably by including members of the symbiont genus *Symbiodinium*.

The nuclear DNA content for 47 dinoflagellate strains is shown in Tables 1 and 2. The DNA content correlated with cell size (Fig. 1; $r^2 = 0.84$ based on an exponential regression trend line, solid gray) and agreed with initial trends observed for dinoflagellates (Rizzo 1987, Veldhuis et al. 1997). The mutualistic endosymbionts of coral reef invertebrates, *Symbiodinium* spp., possessed the smallest genomes (Fig. 1). Several clade B "types," as defined by ITS2 sequence identity (LaJeunesse 2002), had genomes less than $2.0 \text{ pg} \cdot \text{cell}^{-1}$ (Table 2). Therefore, the smallest dinoflagellate genomes were recorded from the smallest *Symbiodinium* spp. (Fig. 1) (LaJeunesse 2001).

The trend illustrated in Figure 1 is similar to most plant and animal cell lineages, with C-values that correlate positively with cell size (Gregory 2001) or carbon content (Holm-Hansen 1969). Numerous explanations for this have been proposed (Gregory 2001), yet it remains unclear whether similar trends exhibited across many diverse and divergent eukaryote lineages are produced by natural selection, historical accident, or simply neutral random processes (Petrov 2001). Because genome size appears to have no relation to genetic and/or organismal complexity ("the DNA C-value paradox," Thomas 1971), it was initially believed that most genomes contained large quantities of secondary "noncoding" DNA representing neutral "junk" (Pagel and Johnstone 1992). Indeed, the DNA content among unicellular dinoflagellates (approximately 1.5 to $170 \text{ pg} \cdot \text{cell}^{-1}$, Tables 1 and 2) is greater than most

TABLE 2. The genome sizes of *Symbiodinium* “types” in culture categorized by clade.

Clade (subgenus)	ITS “type”	Culture I.D. (CCMP no.)	DNA (pg · cell ⁻¹)	Cell size (µm)	Host origin	Geographic origin	
A	A1	61 (2464)	2.2	8–11 (6–7)	<i>Cassiopeia xamachana</i> (Rhizostomeae)	Caribbean (Florida, USA)	
	A1	370 (2467)	2.3		<i>Stylophora pistillata</i> (Scleractinaria)	Red Sea (Gulf of Aqaba)	
	A1.1	80 (2469)	2.3	8–12 (6–7)	<i>Condylactis gigantea</i> (Actiniaria)	Caribbean (Jamaica)	
	A2	185 (2461)	3.4	9–13 (7–8)	<i>Zoanthus sociatus</i>	Caribbean (Jamaica)	
	A3	168 (2457)	3.2		<i>Tridacna crocea</i> (Bivalvia)	South West Pacific (Australia)	
	A3a	PHMS TDle	3.4		<i>Tridacna</i> (Bivalvia)	West Pacific (Phillipines)	
	A4	379 (2456)	2.7		<i>Plexaura homamalla</i> (Gorgonacea)	Caribbean (Bahamas)	
	A12	Culture X	2.2		Unknown	Aquarium reef tank	
	A13	PTA1	2.4		<i>Porites asteroides</i> (Scleractinaria)	Caribbean (Florida, USA)	
	A14	m. miribalis	2.5		<i>Madracis miribalis</i> (Scleractinaria)	Caribbean (Florida, USA)	
	B	B1	2 (2460)	2.1		<i>Aiptasia pallida</i> (Actiniaria)	Caribbean (Florida, USA)
		B1	74	1.9	6–9	<i>Aiptasia pallida</i> (Actiniaria)	Caribbean (Jamaica)
		B1	147 (2470)	1.9	6–8	<i>Pseudotetrogorgia bipinnata</i> (Gorgonacea)	Caribbean (Jamaica)
		B1	Pe	1.9		<i>Pseudotetrogorgia elisabethae</i> (Gorgonacea)	Caribbean (Bahamas)
B2		PPf	2.8		<i>Plexaura flexuosa</i> (Gorgonacea)	Caribbean (Panama)	
B2		10.14b.02	1.5	7.5–11	<i>Montastraea faveolata</i> (Scleractinia)	Caribbean (Florida, USA)	
B2.1		141 (n = 2)	2.5		<i>Oculina diffusa</i> (Scleractinaria)	Western Atlantic (Bermuda)	
B3		385 (2462)	2.4		<i>Dichotomia</i> sp. (Coronatae)	Caribbean	
B19		579	2.6		<i>Briareum</i> sp. newly settled polyp (Gorgonacea)	Caribbean (Florida, USA)	
B25		571	2.4		<i>Briareum</i> sp. newly settled polyp (Gorgonacea)	Caribbean (Florida, USA)	
B26		702	2.9		<i>Plexaura kuna</i> newly settled polyp (Gorgonacea)	Caribbean (Panama)	
C	C1	152 (2466)	4.8	9–13 (6–8)	<i>Rhodactis</i> (<i>Heteractis</i>) <i>lucida</i> (Corallimorph.)	Caribbean (Jamaica)	
	C2	203	2.9		<i>Hippopus hippopus</i> (Bivalvia)	West Pacific (Palau)	
	C2	HHC1B	3.0		<i>Hippopus hippopus</i> (Bivalvia)	West Pacific (Phillipines)	
D	D1a	A001	4.1		<i>Acropora</i> sp. (Scleractinia)	North west Pacific (Okinawa)	
	D1a	2.2b (2556)	3.5	7.5–10	<i>Montastrea faveolata</i> (Scleractinia)	Caribbean (Florida, USA)	
E	E1	383	3.3		<i>Anthopleura sola</i> (= <i>elegantissima</i>) (Actiniaria)	East Pacific (California, USA)	
F	F1 (FR5)	Mv/135 (2468)	3.0	6–11 (4–6)	<i>Montipora verrucosa</i> (Scleractinaria)	Central Pacific (Hawaii, USA)	
	F2 (FR5)	133 (2455)	2.6		<i>Meandrina meandrites</i> (Scleractinaria)	Caribbean (Jamaica)	

The alphanumeric of each ITS “type” refers to the clade (uppercase letter) and ITS sequence identifier (numeral) and, if present, the designation of a co-dominant intragenomic variant (lowercase letter). Example cell size ranges are provided for coccoid and motile phases (in parentheses). The conversion factor for pg to base pair is 1 pg per .965 × 10⁹ bp (Bennett 2000). Center for Culture of Marine Phytoplankton accession numbers are given in parentheses in the Culture I.D. column.

other protist lineages (Velduis et al. 1997) and metazoa (Gregory 2001). However, evidence suggests that DNA content within a nucleus possesses important structural and/or volume-related functions (Cavalier-Smith 1978, Beaton and Cavalier-Smith 1999, Gregory 2001).

Many other dinoflagellate species grow to sizes larger than those we measured. Our survey was limited to smaller isolates for technical reasons: Large and/or nonspherical cells, as they pass through the flow cytometer, can orient in such a way as not to pass through the focus of the illuminations light source and/or of the optics that are used to collect the fluorescence emission. This has the effect of broadening the fluorescence intensity distribution of these types of cells, thereby imparting unacceptable levels of variation on the DNA content measurements. For this reason, species or isolates whose cells were greater than 40 µm in length were not examined.

Has culturing substantially affected genome size? Extended time in culture has been suggested to reduce the C-value and/or increase ploidy in a dinoflagellate strain (Holt and Pfister 1982, Parrow and Burkholder 2002). No indication was found that time in culture had substantially altered the genome sizes of cultured *Symbiodinium* investigated for this study. The C-values measured among different isolates of a particular ITS type were remarkably consistent (Table 2). For, example “A1” cultures #61 originating from *Cassiopea xamachana* in the Caribbean and #370 from *Stylophora pistillata* Esper in the Red Sea differed by 0.1 pg · cell⁻¹ (Table 2). These readings are well within the SE (for technical replications) of our method. Similarly, the values of seven independent measurements of “G.” *simplex* ranged from 9.1 to 12.4 pg · cell⁻¹ with an average of 11.6, variance of 1.53, and SD of 1.24. The “B1” *Symbiodinium* listed in




FIG. 1. Flow cytometric analyses of fluorescence by genomic DNA (pg) stained with DAPI as it relates to cell length (given as the median and range). Increased genome size correlates with increased cell size, a trend observed for most other eukaryotes (Petrov 2001). *Symbiodinium* spp. are among the smallest dinoflagellates, and their genomes are correspondingly small. The representative harmful algal bloom genera *Alexandrium*, *Karenia*, and *Pfiesteria* have 2- to 50-fold larger genomes of *Symbiodinium* spp. Other species listed in Table 1 are represented by black diamonds. An exponential trend line is included ($R^2 = 0.84$). Measurements were made on logarithmically growing cells (Allen et al. 1975).

Table 2 have similar C-values despite being isolated between 5 and 30 years ago from different hosts and geographic regions. Therefore, the close similarity in genome size among various isolates, cultured for various lengths of time, and sharing identical ITS sequences suggests that their genomes are stable in culture (Table 2).

Chromosome volume is proportional to estimated DNA content. The picogram values derived using our DAPI nuclear staining methods are similar to those found using flow-cytometric analysis of fluorescence using SYTOX green (Table 1) (Velduis et al. 1997, Parrow and Burkholder 2002). Both sets of values were proportional to DNA content but typically higher than DNA contents calculated using fluorometric measurements on extracted nucleic acids (Table 1) (Thomas and Farquhar 1978). Although these results are complementary, a second source of evidence supports the accuracy and relative precision of our measurements.

Dinoflagellates and euglenoids are among the few eukaryotic cell types that possess permanently condensed chromosomal bodies (Fig. 2a). The number and size of these is conserved in various species (Holt and Pfister 1982), indicating genetic control. The condensed state therefore allows direct volumetric measurements for each species. Previous calculations made from three-dimensional nuclear reconstructions from various *Symbiodinium* spp. correspond linearly with DAPI fluorescence (Fig. 2b).




FIG. 2. Genome size (DNA content) for described *Symbiodinium* spp. in relation to chromosome volume and number. (a) An electron micrograph through the dinokaryotic nucleus of a *Symbiodinium* sp. showing condensed chromosomes (or "condensed DNA regions," Udy et al. 1993). (b) From left to right, plotted values correspond to the species *S. microadriaticum* subspecies *microadriaticum* (Blank and Huss 1989), *S. microadriaticum* subspecies *condylactis* (Blank and Huss 1989), *S. kawagutii* (Trench and Blank 1987), *S. pilosum* (Trench and Blank 1987), and *S. goreauii* (Trench and Blank 1987). Chromosome volume estimates (μm^3) were based on TEM three-dimensional reconstructions (Trench and Blank 1987, Blank and Huss 1989). Chromosome number and variance are indicated for each taxon.

The accuracy and precision of DNA measurements may be directly attributed to the close similarity in genome size between *Symbiodinium* spp. and the chicken red blood cell standard (Johnston et al. 1999). We observed unusual variability between measurements on the largest species of our survey, *Prorocentrum micans* (225.0 and 115.2 $\text{pg} \cdot \text{cell}^{-1}$ respectively). The wide range in cell size of *P. micans* may explain this variation in DNA concentration (Table 1). Alternatively, because each measurement on this species was taken at separate times, the 2-fold differences in DNA content between dinoflagellate cells in log versus stationary phase (Allen et al. 1975) may better explain this discrepancy.

Endosymbiosis: constraints in the evolution of cell and genome size. Phylogenetic analysis of the dinoflagellates in our survey indicates that genome/cell size has increased and decreased repeatedly over the evolutionary history of this group (Fig. 3). For *Symbiodinium* spp., the ranges in genome and cell sizes are conserved within a narrow range relative to "free-living" dinoflagellate genera (Figs. 1 and 3). The numerous physical, ecological, and evolutionary constraints placed on an intracellular existence may largely explain the conservation of small cell and genome sizes across distantly related *Symbiodinium* spp.

Are *Symbiodinium* genomes smaller than those that would be predicted based on cell size alone? Many pathogenic and mutualistic intracellular bacterial symbionts have smaller genomes than their free-living relatives, often as a result of gene loss and increased




FIG. 3. Maximum likelihood reconstruction based on the 5.8S rRNA gene from genera whose DNA content (in parentheses) are presented in Figure 1. *Symbiodinium* and *Alexandrium* contain taxa with high sequence divergence relative to other genera. A small cell size (6–14 μm) and hence genome size (DNA content) are conserved across various *Symbiodinium* lineages (clade designations are in brackets). This phylogeny, while based on a small number of nucleotides (168 bases), corresponds to previously published reconstructions (Saldarriaga et al. 2001). Numbers associated with internal branches are bootstrap values (>50%) based on 500 replicates. *Perkinsus marinus* was used as the out group taxon (Saldarriaga et al. 2003).

physiological dependence on the host (Moran and Wernegreen 2000). The sister taxa to *Symbiodinium* spp., "*G.*" *simplex* and *Pollarella glacialis* (Montresor et al. 2003), have similar cell lengths (10–12 and 9–14 μm , respectively), yet their genomes are approximately 2.4 to 7 times greater (Fig. 3, Tables 1 and 2). Indeed, values for most *Symbiodinium* spp. fell slightly below the exponential trend line seen in Figure 1. This observation may be an artifact of the size calculations presented for *Symbiodinium* spp. These measurements were based on their coccoid life stage, which tends to be 20%–30% larger than the motile phase (Blank and Huss 1989) (Table 2). Adjustments for this discrepancy would bring *Symbiodinium* spp. C-values more into line with the general trends observed for other dinoflagellates. Although further comparative genomic analyses are needed, the small C-values among *Symbiodinium* spp. may be nothing more than the simple scaling of genome sizes relative to cell size.

Candidates for genome sequencing. It is only a matter of time before sequencing of the first dinoflagellate genome commences. The taxon to be selected should depend on a number of factors, including 1) relative economic and ecological importance, 2) a tractable genome size for sequencing, 3) suitability for genetic manipulation (Lohuis and Miller 1998), and 4) a genome that is representative of the group as a whole. Although *Symbiodinium* clearly meet criteria 1 through 3, unfounded concerns that their associations with invertebrates may have substantially modified their genomes remain. Indeed, they lack many of the complex organelles found in other taxa (Dodge and Greuet 1987). Nonetheless, isolates analyzed in this study are widely distributed host generalists (LaJeunesse 2005) that persist as viable cells in the external environment (Van Oppen et al. 2001, LaJeunesse et al. 2004). They are easily isolated from the host and maintained "happily" in culture as independent free-living dinoflagellates (host "specialists," however, have yet to be cultured, personal observation). Therefore, these are typical dinoflagellates not lacking important housekeeping genes. Given all these factors and considerations, we believe *Symbiodinium* spp. represent leading candidates for genome sequencing.

We thank Robert Moore for pointing out to us the need for additional genome measurements including the various *Symbiodinium* spp. R. K. Trench, D. Poland, N. Kirk, and J. Holmberg contributed comments on the manuscript. Funding for TCL was provided by an NSF grant (OCE-0137007) and FIU start-up. M. A. C.'s contributions were supported by NSF grants OCE-99-07319 and NSF OCE 03-13708 and R. A. A.'s contributions were supported by NSF grant DBI-9910676. C. Lewis provided technical assistance, and the Niagara Falls Aquarium, Niagara Falls, New York supplied seawater for maintaining algal cultures at the SUNY Buffalo laboratory.

Allen, J. R., Roberts, M., Loeblich III, A. R. & Klotz, L. C. 1975. Characterization of the DNA from the dinoflagellate *Cryptohcodinium cohnii* and implications for nuclear organization. *Cell* 6:161–9.

Baker, A. C. 2003. Flexibility and specificity in coral-algal symbiosis: diversity, ecology and biogeography of *Symbiodinium*. *Annu. Rev. Ecol. Syst.* 34:661–89.

Beaton, M. J. & Cavalier-Smith, T. 1999. Eukaryotic non-coding DNA is functional: evidence from the differential scaling of cryptomonad genomes. *Proc. R. Soc. Lond.* 266:2053–9.

Bennett, M. D. 2000. Variation in genomic form in plants and its ecological implications. *New Phytol.* 106:177–200.

Blank, R. J. 1987. Cell architecture of the dinoflagellate *Symbiodinium* sp. inhabiting the Hawaiian stony coral *Montipora verrucosa*. *Mar. Biol.* 94:143–55.

Blank, R. J. & Huss, V. A. R. 1989. DNA divergency and speciation in *Symbiodinium* (Dinophyceae). *Pl. Syst. Evol.* 163:153–63.

Blank, R. J. & Trench, R. K. 1985. Speciation and symbiotic dinoflagellates. *Science* 229:656–8.

Cachon, J. M. 1987. Parasitic dinoflagellates. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 571–610.

Cavalier-Smith, T. 1978. Nuclear volume control by nucleoskeletal DNA, selection for cell volume and cell growth rate, and the solution of the DNA C-value paradox. *J. Cell Sci.* 34:247–78.

Coats, D. W. 2002. Dinoflagellate life-cycle complexities. *J. Phycol.* 38:417–9.

- Daugbjerg, N., Hansen, G., Larsen, J. & Moestrup, Ø. 2000. Phylogeny of some of the major genera of dinoflagellates based on ultrastructure and partial LSU rDNA sequence data, including the erection of three new genera of unarmoured dinoflagellates. *Phycologia* 39:302–17.
- Dodge, J. D. & Greuet, C. 1987. Dinoflagellate ultrastructure and complex organelles. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 92–142.
- Domotor, S. L. & D'Elia, C. F. 1986. Cell-size distributions of zooxanthellae in culture and symbiosis. *Biol. Bull.* 170:519–25.
- Elela, S. A. & Nazar, R. N. 1997. Role of the 5.8S rRNA in ribosomal translocation. *Nucleic Acids Res.* 25:1788–94.
- Fitt, W. K. 1985. Effect of different strains of the zooxanthella *Symbiodinium microadriaticum* on growth and survival of their coelenterate and molluscan hosts. *Proc. 5th Int. Coral Reef Congr.* 6:131–6.
- Gregory, T. R. 2001. Coincidence, coevolution, or causation? DNA content, cell size, and the C-value enigma. *Biol. Rev. Camb. Phil. Soc.* 76:65–101.
- Holm-Hansen, O. 1969. Algae: amounts of DNA and organic carbon in single cells. *Science* 163:87–8.
- Holt, J. R. & Pfister, L. A. 1982. A technique for counting chromosomes of armored dinoflagellates, and chromosome numbers of six freshwater dinoflagellate species. *Am. J. Bot.* 69:1165–8.
- Johnston, J. S., Bennett, M. D., Rayburn, A. L., Galbraith, D. W. & Price, J. H. 1999. Reference standards for determination of DNA content of plant nuclei. *Am. J. Bot.* 86:609–13.
- Lajeunesse, T. C. 2001. Investigating the biodiversity, ecology, and phylogeny of endosymbiotic dinoflagellates in the genus *Symbiodinium* using the internal transcribed spacer region: in search of a "species" level marker. *J. Phycol.* 37:866–80.
- Lajeunesse, T. C. 2002. Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Mar. Biol.* 141:387–400.
- Lajeunesse, T. C. 2005. "Species" radiations of symbiotic dinoflagellates in the Atlantic and Indo-Pacific since the Miocene-Pliocene transition. *Mol. Biol. Evol.* 22:570–81.
- Lajeunesse, T. C., Bhagooli, R., Hidaka, M., Done, T., deVantier, L., Schmidt, G. W., Fitt, W. K. & Hoegh-Guldberg, O. 2004. Closely-related *Symbiodinium* spp. differ in relative dominance within coral reef host communities across environmental, latitudinal, and biogeographic gradients. *Mar. Ecol. Prog. Ser.* 284:147–61.
- Loeblich III, A. R. 1976. Dinoflagellate genetics and DNA characterization. *Stadler Symp.* 8:111–28.
- Lohuis, M. R. ten & Miller, D. J. 1998. Genetic transformation of dinoflagellates (*Amphidinium* and *Symbiodinium*): expression of GUS in microalgae using heterologous promoter constructs. *Plant J.* 13:427–35.
- Maroteaux, L., Herzog, M. & Soyer-Gobillard, M. 1985. Molecular organization of dinoflagellate ribosomal DNA: evolutionary implications of the deduced 5.8 S rRNA secondary structure. *Biosystems* 18:307–19.
- Montresor, M., Lovejoy, C., Orisini, L., Procaccini, G. & Roy, S. 2003. Bipolar distribution of the cyst-forming dinoflagellate *Polarella glacialis*. *Polar Biol.* 26:186–94.
- Moran, A. N. & Wernegreen, J. J. 2000. Are mutualism and parasitism irreversible evolutionary alternatives for endosymbiotic bacteria? Insights from molecular phylogenetics and genomics. *Trends Ecol. Evol.* 15:321–6.
- Muscatine, L. 1990. The role of symbiotic algae in carbon and energy flux in reef corals. In Dubinsky, Z. [Ed.] *Ecosystems of the World, Vol. 25. Coral Reefs*. Elsevier, Amsterdam, pp. 75–87.
- Muscatine, L. & Porter, J. W. 1977. Reef corals: mutualistic symbioses adapted to nutrient-poor environments. *BioScience* 27:454–60.
- Pagel, M. & Johnstone, R. A. 1992. Variation across species in the size of the nuclear genome supports the junk-DNA explanation for the C-value paradox. *Proc. R. Soc. Lond. B* 249:119–24.
- Parrow, M. W. & Burkholder, J. M. 2002. Flow cytometric determination of zoospore DNA content and population DNA distribution in cultured *Pfiesteria* spp. (Pyrrhophyta). *J. Exp. Mar. Biol. Ecol.* 267:35–51.
- Petrov, D. A. 2001. Evolution of genome size: new approaches to an old problem. *Trends Genet.* 17:23–8.
- Pfister, L. A. & Anderson, D. M. 1987. Dinoflagellate reproduction. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 611–48.
- Rizzo, P. J. 1987. Biochemistry of the dinoflagellate nucleus. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 143–73.
- Saldarriaga, J. F., McEwan, M. L., Fast, N. M., Taylor, F. J. R. & Keeling, P. J. 2003. Multiple protein phylogenies show that *Oxyrrhis marina* and *Perkinsus marinus* are early branches of the dinoflagellate lineage. *Int. J. Syst. Evol. Micro.* 53:355–65.
- Saldarriaga, J. F., Taylor, F. J. R., Keeling, P. J. & Cavalier-Smith, T. 2001. Dinoflagellate nuclear SSU rRNA phylogeny suggests multiple plastid losses and replacements. *J. Mol. Evol.* 53:204–13.
- Santos, S. R. & Coffroth, M. A. 2003. Molecular genetic evidence that dinoflagellates belonging to the genus *Symbiodinium* Freudenthal are haploid. *Biol. Bull.* 204:10–20.
- Schoenberg, D. A. & Trench, R. K. 1980. Genetic variation in *Symbiodinium* (= *Gymnodinium*) *microadriaticum* Freudenthal, and specificity in its symbiosis with marine invertebrates. II. Morphological variation in *S. microadriaticum*. *Proc. R. Soc. Lond. B* 207:429–44.
- Shimizu, Y. 1987. Dinoflagellate toxins. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 282–315.
- Swofford, D. L. 2000. *Paup*. Phylogenetic Analysis Using Parsimony (*And Other Methods)*. Version 4.0b10. Sinauer, Sunderland, MA.
- Taylor, F. J. R. [Ed.] 1987. *The Biology of Dinoflagellates*. Blackwell, Oxford, 785 pp.
- Thomas, C. A. 1971. The generic organization of chromosomes. *A. Rev. Genet.* 5:237–56.
- Thomas, P. S. & Farquhar, M. N. 1978. Specific measurement of DNA in nuclei and nucleic acids using diaminobenzoic acid. *Anal. Biochem.* 89:35–44.
- Trench, R. K. 1993. Microalgal-invertebrate symbioses: a review. *Endocytobiosis Cell Res.* 9:135–75.
- Trench, R. K. & Blank, R. J. 1987. *Symbiodinium microadriaticum* Freudenthal, *S. goreauii* sp. nov., *S. kawagutii* sp. nov. and *S. pilosum* sp. nov.: gymnodinoid dinoflagellate symbionts of marine invertebrates. *J. Phycol.* 23:469–81.
- Triplett, E. L., Govind, N. S., Roman, S. J., Jovine, R. V. M. & Prezelin, B. B. 1993. Characterization of the sequence organization of DNA from the dinoflagellate *Heterocapsa pygmaea* (*Glenodinium* sp.). *Mol. Mar. Bio. Biotech.* 2:239–45.
- Udy, J. W., Hinde, R. & Vesik, M. 1993. Chromosomes and DNA in *Symbiodinium* from Australian hosts. *J. Phycol.* 29:314–20.
- Van Oppen, M. J. H., Palstra, F. P., Piquet, A. M.-T. & Miller, D. J. 2001. Patterns of coral-dinoflagellate associations in *Acropora*: significance of local availability and physiology of *Symbiodinium* strains and host-symbiont selectivity. *Proc. R. Soc. Lond. B* 268:1759–67.
- Veldhuis, M. J. W., Cucci, T. L. & Sieracki, M. E. 1997. Cellular DNA content of marine phytoplankton using two new fluorochromes: taxonomic and ecological implications. *J. Phycol.* 33:527–41.
- Whitney, S. M., Shaw, D. C. & Yellowlees, D. 1995. Evidence that some dinoflagellates contain a ribulose-1,5-bisphosphate carboxylase/oxygenase related to that of the α -proteobacteria. *Proc. R. Soc. Lond. B* 259:271–5.
- Withers, N. W. 1987. Dinoflagellate sterols. In Taylor, F. J. R. [Ed.] *The Biology of Dinoflagellates*. Blackwell, Oxford, pp. 316–59.
- Zhang, Z., Green, B. R. & Cavalier-Smith, T. 1999. Single gene circles in dinoflagellate chloroplast genomes. *Nature* 400:155–9.