

Menthol-induced bleaching rapidly and effectively provides experimental aposymbiotic sea anemones (*Aiptasia* sp.) for symbiosis investigations

Jennifer L. Matthews¹, Ashley E. Sproles¹, Clinton A. Oakley¹, Arthur R. Grossman², Virginia M. Weis³ and Simon K. Davy*¹

¹School of Biological Sciences, Victoria University of Wellington, Wellington 6140, New Zealand

²Department of Plant Biology, The Carnegie Institution, Stanford, California 94305, United States

³Oregon State University, Department of Integrative Biology, 3029 Cordley Hall, Corvallis, OR 97331, United States

*Corresponding author: Simon.Davy@vuw.ac.nz

Key words: Cnidarian-dinoflagellate symbiosis; *Symbiodinium*; Coral reefs

Abstract

Experimental manipulation of the symbiosis between cnidarians and photosynthetic dinoflagellates (*Symbiodinium* spp.) is critical to advance understanding of the cellular mechanisms involved in host–symbiont interactions, and overall coral reef ecology. The anemone *Aiptasia* sp. is a model for the cnidarian-dinoflagellate symbiosis, and notably it can be rendered aposymbiotic (i.e. dinoflagellate-free) and re-infected with a range of *Symbiodinium* types. Various methods exist for generating aposymbiotic hosts, however they can be hugely time-consuming and not wholly effective. Here, we optimise a method using menthol for production of aposymbiotic *Aiptasia*. The menthol treatment produced aposymbiotic hosts within just four weeks (97–100% symbiont loss), which was maintained long after treatment when anemones were held under a standard light/dark cycle. The ability of *Aiptasia* to form a stable symbiosis appeared unaffected by menthol exposure, as demonstrated by successful symbiosis re-establishment when experimentally re-infected. Furthermore, there was no significant impact on photosynthetic or respiratory performance of re-infected anemones.

Introduction

The symbiosis between corals and dinoflagellate algae of the genus *Symbiodinium* promotes the success of coral reefs in nutrient-poor tropical seas. However, there are still large gaps in our knowledge about how this symbiosis is established and maintained (Davy et al., 2012; Weis and Allemand, 2009; Weis et al., 2008). Moreover, we do not fully understand how and why this symbiosis breaks down under stress (i.e. bleaching) (Weis, 2008). These knowledge gaps hinder our capacity to understand the function of coral reefs, and to predict how coral reefs might respond to our changing environment.

In order to elucidate the cellular basis of the coral-dinoflagellate relationship, the symbiotic sea anemone *Aiptasia* sp. has been widely adopted as a model system due to its robust nature and the ability to produce large populations easily in the laboratory (Weis et al., 2008). Furthermore, *Aiptasia* sp. can be cleared of symbionts, maintained

in an aposymbiotic state for extended periods, and successfully re-infected with a variety of *Symbiodinium* types (Belda-Baillie et al., 2002; Schoenberg and Trench, 1980; Starzak et al., 2014). The ease of experimental manipulation combined with the rapidly increasing resolution of genomic, proteomic and metabolic databases for *Aiptasia* sp., makes this symbiotic anthozoan an ideal candidate for biochemical and genetic experiments that will further our understanding of the cellular processes underlying the cnidarian-dinoflagellate symbiosis (Lehnert et al., 2012; Peng et al., 2010; Sunagawa et al., 2009; Weis et al., 2008).

Currently, the most commonly used method to render anemones (including *Aiptasia* sp.) aposymbiotic, combines temperature stress (heat- or cold-shock), followed by dark treatment and/or chemical inhibition of photosynthesis with 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) (e.g. Belda-Baillie et al., 2002; Lehnert et al., 2014; Xiang et al., 2013). This is a slow, laborious process which often requires months of preparation (Starzak et al., 2014; Xiang et al., 2013), and may not result in the full eradication of *in hospite* endosymbionts (Belda-Baillie et al., 2002; Schoenberg and Trench, 1980; Wang and Douglas, 1998).

Recently, Wang et al. (2012) described a method of menthol-induced bleaching in the corals *Isopora palifera* and *Stylophora pistillata*. Menthol is known to cause local anesthetic effects in neuronal and skeletal muscles *via* blocking voltage-operated sodium channels (Haeseler et al., 2002). This cellular response has led to its use as a marine anaesthetic (Alexander, 1964; Laretta et al., 2014). Menthol is known to act on a variety of different membrane receptors, including transient receptor potential (TRP)M8 that results in an increase in intracellular Ca^{2+} concentrations and causes a cold sensation in vertebrates (Hans et al., 2012; McKemy et al., 2002; Okazawa et al., 2000; Peier et al., 2002). As proposed by Wang et al. (2012), the mechanism of menthol-induced bleaching might be attributable to Ca^{2+} stimulated exocytosis (Pang and Südhof, 2010). Preliminary experiments conducted by Wang et al. (2012) also suggest that menthol inhibition of *Symbiodinium* photosystem II activity may play a role in the expulsion of the algal cells or the digestion of the *Symbiodinium* cells by the host. Although the exact mechanism by which menthol induces symbiont

expulsion is not yet clear, the findings of Wang and co-authors provide a platform for developing the use of menthol to generate aposymbiotic *Aiptasia* sp.

We therefore tested the applicability of this approach to *Aiptasia* sp., in terms of its effectiveness and impact on the capacity for symbiosis reestablishment. In particular, we determined whether menthol can rapidly and effectively produce aposymbiotic anemones, and whether these anemones can be experimentally re-infected with symbiotic dinoflagellates to similar cell densities as untreated and healthy symbiotic anemones. Ultimately, we describe a method to assist the many researchers around the world who use the *Aiptasia* model system to further our understanding of the coral-dinoflagellate symbiosis.

Materials and Methods

Experimental organisms

Symbiotic individuals of *Aiptasia* sp. ($n = 300$) were harvested from a long-term lab stock maintained in 1 μm filtered seawater (FSW) at 25°C, with light provided by AQUA-GLO T8 fluorescent bulbs at $\sim 95 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ (light:dark = 12 h:12 h). Anemones were maintained in 1 μm filtered seawater (FSW) aerated with a Hailea® aquarium air pump at 2 L/min and at 25°C using a WEIPRO® Temperature Controller MX-1019. These lighting and aquarium conditions were maintained throughout the 19-week experiment. The anemones were evenly divided among the treatments (menthol, cold shock, and untreated) and allowed to settle for 72 h before beginning a 4-week treatment period.

The *Symbiodinium* culture (ID: FLAp2) was grown at 25°C and an irradiance of 100 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ on a 12:12 h light:dark cycle. *Symbiodinium* cells were sub-cultured from a laboratory stock and grown in silica-free f2 medium (AusAqua PtyLtd, SA, Australia) for six weeks before use in the infection study. To genotype the *Symbiodinium* cells, DNA was extracted as described by Hill et al. (2014) from an algal pellet derived from a 1 ml aliquot of the *Symbiodinium* suspension. Samples

were immediately used for PCR, using the thermal cycling regime and reaction mixture of Hill et al. (2014), and the outer primers ITSintfor2 and ITS2Rev2. PCR products were cleaned with ExoSAP-IT (USB Corporation, OH, USA) and sequenced by the Macrogen Sequencing Service (Macrogen Inc., Seoul, South Korea). Sequences were aligned with Geneious v. 7.0 (Biomatters Ltd., Auckland, NZ) and a BLAST search carried out against *Symbiodinium* ITS2 sequences in GenBank. Using this approach, *Symbiodinium* genotype was confirmed as B1.

Menthol versus cold-shock treatment

Menthol (20% w/v in ethanol; Sigma-Aldrich, Auckland, NZ) was added to 1 μm FSW at a final concentration of 0.19 mM. This concentration resulted in successful bleaching without causing mortality. A higher concentration of 0.38 mM as suggested by Wang et al. (2012) was trialled prior to this experiment, but caused mortality of over 50% of the anemones (unpublished data). The anemones were incubated in the menthol/FSW solution for 8 h after which the menthol/FSW was removed and the anemones incubated in 1 μm FSW for 16 h. DCMU (100 mM dissolved in EtOH, Sigma-Aldrich, Auckland, NZ) was added to a final concentration of 5 μM , in order to prevent nuisance algal blooms and to limit the re-establishment of any residual symbiont cells by inhibiting photosynthesis. This 24-h cycle was repeated for four consecutive days. The 8-h menthol/FSW incubation occurred during the 12-h light period. The 4-day treatment was repeated after a 3-day break, during which anemones were maintained under the lighting and temperature conditions stated above, with biweekly feeding with *Artemia* sp. nauplii.

For cold shock, the water was replaced with 4°C 1 μm -FSW, and placed in a refrigerator at 4°C for 4 h. Afterwards, the water was replaced with 25°C 1 μm -FSW containing a final concentration of 50 μM DCMU (100 mM dissolved in EtOH, Sigma-Aldrich, Auckland, NZ) and held under the lighting and aquarium conditions described above. The anemones were incubated under these conditions for three days, and the treatment then repeated. The anemones were unfed for the duration of the cold-shock, as *per* the protocol described by Steen and Muscatine (1987). Anemones were cold-shocked a total of eight times (twice *per* week) during the treatment period.

After the 4-week treatment period, all anemones were maintained in 1 μm -FSW at 25°C under the irradiance regime described above, with biweekly feeding, for an additional 15 weeks. Their symbiotic status was assessed *via* confocal fluorescence microscopy and cell density estimates (described below) at Weeks 1-5, 7, 9, 11, and 19.

The untreated group of anemones was kept under the same thermal and light regimes as the menthol-treated and cold-shocked anemones, with biweekly feeding throughout the entire 19-week experiment. They were assessed for symbiont density at the same time-points as the experimental anemones, to obtain an average symbiont density of untreated, healthy anemones.

Reinfection of aposymbiotic anemones

The capacity to re-infect menthol-treated anemones with algal symbionts was tested; this was not possible with cold-shocked anemones since they were not fully aposymbiotic. Anemones were infected with cultured homologous *Symbiodinium* type B1, *i.e.* symbionts originally isolated from *Aiptasia* sp. A drop of concentrated *Symbiodinium* suspension (3×10^6 cells ml^{-1}) was pipetted onto the oral disc of each anemone, followed by a drop of *Artemia* sp. nauplii to invoke a feeding response and encourage *Symbiodinium* uptake (Davy et al. 1997). This process was repeated 4 h later, and then the water was replaced with 1 μm -FSW 8 h after the second infection.

Confocal microscopy

Whole anemones from each treatment were examined at the same time-points as cell density assessments by confocal microscopy (Olympus Provis AX70, at 100 \times magnification) to detect the chlorophyll autofluorescence of any dinoflagellates. Anemones were selected at random from each treatment, and placed in a “relaxation solution” [50% FSW, 50% 0.37 M magnesium chloride (MgCl_2)] for 15 min before being moved to a FluoroDishTM glass-bottom confocal dish (World Precision Instruments, Florida USA). The auto-fluorescence emission of the *Symbiodinium* cells

was excited using a 559 nm laser and captured at $647 \text{ nm} \pm 10 \text{ nm}$. This was performed to check the rate of bleaching and the rate of reinfection, as well as to detect any background *Symbiodinium* populations in the post-treatment anemones.

Symbiont density and protein content

Symbiodinium populations were quantified with a haemocytometer and light microscope ($100 \times$ magnification; $n = 8$ replicate counts *per* sample) using whole anemone homogenate ($n = 5$). Cell density was normalised to soluble protein content, measured *via* the Bradford assay (Bradford, 1976). Cell density was measured at Weeks 1–5, 7, 9, 11, and 19.

O₂ flux

Fifteen weeks after reinfection, maximum photosynthetic and dark respiratory O₂ fluxes were measured to compare re-infected menthol-treated anemones against untreated anemones ($n = 4$). A FIBOX 3 fibre optic oxygen transmitter and oxygen probe (PreSens GmbH, Germany) were used as described by Starzak et al. (2014). The respiration rate ($\text{mL O}_2 \text{ h}^{-1}$) was measured for 30 min in the dark, after which time the rate of net photosynthesis was measured for 30 min at an irradiance of $\sim 100 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. Net photosynthesis and respiration were corrected for background levels using a FSW-only control, and gross photosynthesis was calculated by the addition of net photosynthesis to dark respiration (Muscatine et al. 1981). The average ratio of gross photosynthesis to respiration (P:R) for 24 hours was calculated for each anemone, by assuming light:dark = 12 h:12 h and that respiration was constant in the light and dark, and the average ratio for each treatment calculated.

Statistical tests

The nonparametric Mann-Whitney U test was used to test for differences between the treatments with respect to symbiont density (SPSS statistical software, v. 20, IBM Corp). Separate analyses were run for the bleaching and post-bleaching periods.

A one-way analysis of variance (ANOVA) was performed on the photosynthetic rate, respiration rate, and log-transformed P:R values to determine whether significant differences were present between the re-infected menthol-treated and untreated symbiotic anemones at the end of the 19-week experimental period.

Results and Discussion

Menthol treatment induced 97–100% symbiont loss within 14–28 days, with 60% of anemones losing all symbionts. In comparison, cold shock never induced 100% loss within the same time frame (Fig. 1A, B), and any residual symbionts in the menthol-treated anemones did not re-establish a full symbiosis once returned to normal seawater conditions over the course of our experiment (Fig. 1A, C).

While statistically, the patterns of symbiont loss during Weeks 1-4 were not different between the menthol and cold-shock treatments (Fig. 1A; Mann-Whitney $U = 113$, $df = 4$, $P = 0.423$, $n = 5$), only the menthol-treated anemones successfully maintained aposymbiotic status in the longer term. In both treatments, symbiont density remained near zero during the first week of recovery (Weeks 4-5; Fig. 1A), however subsequently there was a significant difference between the cell densities of menthol- and cold shock-treated anemones (Mann-Whitney $U = 35$, $df = 4$, $P < 0.0005$; $n = 5$), with the menthol-treated anemones remaining mostly symbiont-free (0-0.05% of pre-bleaching density) and the cold-shocked anemones becoming fully re-populated (Weeks 5-9; Fig. 1A). Confocal microscopy confirmed these measurements (Fig. 1C). Menthol treatment therefore has considerable potential for the rapid and effective generation of aposymbiotic *Aiptasia* sp.

One key use for aposymbiotic *Aiptasia* sp. is for experimental work that involves re-infection with symbiotic algae of different types and from different host species. It was therefore important to test the infectivity of the menthol-treated anemones, and establish whether there were any significant long-term impacts of menthol treatment *versus* no treatment on physiological performance (photosynthesis and respiration) of the re-infected anemones. When infected with *Symbiodinium* cells (ITS2 type B1), symbiont density increased slowly during the initial two weeks, followed by a rapid

increase during Weeks 2–12 (Fig. 2A). Confocal microscopy confirmed that menthol-treated anemones were fully repopulated after 12 weeks (Fig. 2B). The infection pattern and maximum cell density were similar to other re-infection studies using heat- and cold-shock bleached *Aiptasia* and type B1 *Symbiodinium*, where a steep increase in cell density was observed in the first four weeks, followed by stabilization (Belda-Baillie et al. 2002; Schoenberg and Trench, 1980; Starzak et al. 2014). Additionally, the photosynthetic rate ($P = 0.432$; $n = 4$), respiration rate ($P = 0.347$; $n = 4$), and ratio of photosynthesis to respiration ($P = 0.368$; $n = 4$) in these anemones were similar to those of untreated symbiotic anemones. This indicates that their capacity to re-form a functional symbiosis was not compromised, further highlighting the value of this method.

As expected given its anaesthetic properties (Alexander, 1964; Laretta et al., 2014), menthol induced tentacle relaxation and unresponsiveness in anemones for the duration of the treatment. Once the seawater was replaced with regular FSW, the anemones were able to regain full movement and feeding capabilities within ~15 min. This is in stark contrast to the residual effects of cold shock on anemones, where long-lasting tentacle retraction and occasional mortality were observed.

In summary, we provide a rapid and effective method for the generation of aposymbiotic *Aiptasia* sp. that will facilitate studies of symbiosis establishment and maintenance, host-symbiont recognition, and symbiosis function with this model system. In particular, menthol treatment has the potential to rapidly create large stocks of aposymbiotic anemones without compromising their infectivity. Ultimately, this method will help to accelerate the rate of discovery as we try to better understand the function of coral reefs, and their response and adaptive capacity in the face of climate change.

List of Symbols and Abbreviations

DCMU – 3-(3,4-dichlorophenyl)-1,1-dimethylurea

FSW – filtered seawater

Competing interests

No competing interests declared.

Author contributions

J.L.M., A.E.S., C.A.O., A.R.G., V.M.W. and S.K.D. conceived and designed the experiments. J.L.M. and A.E.S. carried out the experiments and subsequent data analysis. J.L.M., A.E.S., C.A.O., A.R.G., V.M.W. and S.K.D. wrote the manuscript.

Funding

This work was supported by a grant from the Royal Society of New Zealand Marsden Fund (Grant number 1202; awarded to S.K.D., A.R.G and V.M.W.).

References

- Alexander, R. M.** (1964). Visco-elastic properties of the mesogloea of jellyfish. *Journal of Experimental Biology* **41**, 363-369.
- Belda-Baillie, C. A., Baillie, B. K. and Maruyama, T.** (2002). Specificity of a model cnidarian-dinoflagellate symbiosis. *The Biological Bulletin* **202**, 74-85.
- Brown, B.** (1997). Coral bleaching: causes and consequences. *Coral Reefs* **16**, S129-S138.
- Davy, S. K., Allemand, D. and Weis, V. M.** (2012). Cell biology of cnidarian-dinoflagellate symbiosis. *Microbial and Molecular Biology Reviews* **76**, 229-61.
- Haeseler, G., Maue, D., Grosskreutz, J., Bufler, J., Nentwig, B., Piepenbrock, S., Dengler, R. and Leuwer, M.** (2002). Voltage-dependent block of neuronal and skeletal muscle sodium channels by thymol and menthol. *European Journal of Anaesthesiology* **19**, 571-579.
- Hans, M., Wilhelm, M. and Swandulla, D.** (2012). Menthol suppresses nicotinic acetylcholine receptor functioning in sensory neurons via allosteric modulation. *Chemical Senses* **37**, 463-469.
- Hill, R., Fernance, C., Wilkinson, S. P., Davy, S. K., Scott, A.** (2014). Symbiont shuffling during thermal bleaching and recovery in the sea anemone *Entacmaea quadricolor*. *Marine Biology* **161**, 2931-2937
- Lauretta, D., Häussermann, V., Brugler, M. R. and Rodríguez, E.** (2014). *Isoparactis fionae* sp. nov.(Cnidaria: Anthozoa: Actiniaria) from Southern Patagonia with a discussion of the family Isanthidae. *Organisms Diversity & Evolution* **14**, 31-42.
- Lehnert, E. M., Burriesci, M. S. and Pringle, J. R.** (2012). Developing the anemone *Aiptasia* as a tractable model for cnidarian-dinoflagellate symbiosis: the transcriptome of aposymbiotic *A. pallida*. *BMC Genomics* **13**, 271.
- Lehnert, E. M., Mouchka, M. E., Burriesci, M. S., Gallo, N. D., Schwarz, J. A. and Pringle, J. R.** (2014). Extensive differences in gene expression between symbiotic and aposymbiotic cnidarians. *G3 (Bethesda)* **4**, 277-95.
- Little, A. F., van Oppen, M. J. and Willis, B. L.** (2004). Flexibility in algal endosymbioses shapes growth in reef corals. *Science* **304**, 1492-4.

- McKemy, D. D., Neuhauser, W. M. and Julius, D.** (2002). Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* **416**, 52-58.
- Muscatine, L., Grossman, D. and Doino, J.** (1991). Release of symbiotic algae by tropical sea anemones and corals after cold shock. *Marine Ecological Series* **20**, 233-243
- Okazawa, M., Terauchi, T., Shiraki, T., Matsumura, K. and Kobayashi, S.** (2000). 1-Menthol-induced $[Ca^{2+}]_i$ increase and impulses in cultured sensory neurons. *Neuroreport* **11**, 2151-2155.
- Pang, Z. P. and Südhof, T. C.** (2010). Cell biology of Ca^{2+} -triggered exocytosis. *Current Opinion in Cell Biology* **22**, 496-505.
- Peier, A. M., Moqrich, A., Hergarden, A. C., Reeve, A. J., Andersson, D. A., Story, G. M., Earley, T. J., Dragoni, I., McIntyre, P. and Bevan, S.** (2002). A TRP channel that senses cold stimuli and menthol. *Cell* **108**, 705-715.
- Peng, S. E., Wang, Y. B., Wang, L. H., Chen, W. N. U., Lu, C. Y., Fang, L. S. and Chen, C. S.** (2010). Proteomic analysis of symbiosome membranes in Cnidaria–dinoflagellate endosymbiosis. *Proteomics* **10**, 1002-1016.
- Sawyer, S. J. and Muscatine, L.** (2001). Cellular mechanisms underlying temperature-induced bleaching in the tropical sea anemone *Aiptasia pulchella*. *Journal of Experimental Biology* **204**, 3443-3456.
- Schoenberg, D. and Trench, R.** (1980). Genetic variation in *Symbiodinium* (= *Gymnodinium*) *microadriaticum* Freudenthal, and specificity in its symbiosis with marine invertebrates. III. Specificity and Infectivity of *Symbiodinium microadriaticum*. *Royal Society of London Proceedings Series B* **207**, 445-460
- Starzak, D. E., Quinnell, R. G., Nitschke, M. R. and Davy, S. K.** (2014). The influence of symbiont type on photosynthetic carbon flux in a model cnidarian–dinoflagellate symbiosis. *Marine Biology* **161**, 711-724.
- Steen, R. G. and Muscatine, L.** (1987). Low temperature evokes rapid exocytosis of symbiotic algae by a sea anemone. *The Biological Bulletin* **172**, 246-263.
- Sunagawa, S., Wilson, E. C., Thaler, M., Smith, M. L., Caruso, C., Pringle, J. R., Weis, V. M., Medina, M. and Schwarz, J. A.** (2009). Generation and analysis of transcriptomic resources for a model system on the rise: the sea anemone *Aiptasia pallida* and its dinoflagellate endosymbiont. *BMC Genomics* **10**, 258.

- Wang, J. and Douglas, A.** (1998). Nitrogen recycling or nitrogen conservation in an alga-invertebrate symbiosis? *Journal of Experimental Biology* **201**, 2445-2453.
- Weis, V. M.** (2008). Cellular mechanisms of Cnidarian bleaching: stress causes the collapse of symbiosis. *Journal of Experimental Biology* **211**, 3059-66.
- Weis, V. M. and Allemand, D.** (2009). Physiology. What determines coral health? *Science* **324**, 1153-5.
- Weis, V. M., Davy, S. K., Hoegh-Guldberg, O., Rodriguez-Lanetty, M. and Pringle, J. R.** (2008). Cell biology in model systems as the key to understanding corals. *Trends in Ecology and Evolution* **23**, 369-76.
- Xiang, T., Hambleton, E. A., DeNofrio, J. C., Pringle, J. R., Grossman, A. R. and Lin, S.** (2013). Isolation of clonal axenic strains of the symbiotic dinoflagellate *Symbiodinium* and their growth and host specificity. *Journal of Phycology* **49**, 447-458.

Figures




Figure 1. Effects of menthol and cold shock on algal cell density in *Aiptasia* sp. (A) Density of *Symbiodinium* cells in menthol-treated (closed circles) and cold-shocked (open circles) anemones over a 19-week period. The anemones were treated for the first four weeks, then left to recover in regular FSW for the remaining 15 weeks. Outliers (defined as exceeding $1.5 \times$ IQR) were detected in the menthol-treated anemones at Weeks 4 and 11, and are indicated by a red cross. Values are mean \pm s.e.m.; $n = 5$. (B) Confocal microscopy images of anemones during menthol treatment and cold shock (Weeks 1-4). Yellow arrows indicate *Symbiodinium* cells in

host tissues. **(C & D)** Confocal fluorescence images of menthol-treated and cold-shocked anemones, respectively, 15 weeks post-treatment (i.e. Week 19). Scale bars are 100 μm .




Figure 2. Repopulation of menthol-treated *Aiptasia* sp. with algal symbionts. (A) Symbiont cell density over a 12-week period after anemones were experimentally re-infected with the homologous *Symbiodinium* type B1 (closed triangles). Also shown are symbiont densities in symbiotic, untreated anemones (closed square) and menthol-treated anemones that were not experimentally re-infected (open circles). Values are mean \pm s.e.m.; $n = 5$. (B & C) Confocal images showing, respectively, the re-established symbiont population 12 weeks after reinfection *versus* tissues in a non-infected aposymbiotic anemone. Scale bars are 100 μ m.